

10594448

File 5:Biosis Previews(R) 1926-2009/Feb W1
(c) 2009 The Thomson Corporation

Set	Items	Description
? s p73 and IKK?	1565	P73
	2942	IKK?
S1	2	P73 AND IKK?

? t s1/7/1-2

1/7/1

DIALOG(R)File 5:Biosis Previews(R)
(c) 2009 The Thomson Corporation. All rts. reserv.

0020183549 BIOSIS NO.: 200800230488

ATM-dependent nuclear accumulation of $\text{IKK}\alpha$ plays an important role in the regulation of p73 -mediated apoptosis in response to cisplatin

AUTHOR: Yoshida K; Ozaki T; Furuya K; Nakanishi M; Kikuchi H; Yamamoto H; Ono S; Koda T; Omura K; Nakagawara A (Reprint)

AUTHOR ADDRESS: Chiba Canc Ctr Res Inst, Div Biochem, Chuoh Ku, 666-2 Nitona, Chiba 2608717, Japan**Japan

AUTHOR E-MAIL ADDRESS: akiranak@chiba-cc.jp

JOURNAL: Oncogene 27 (8): p1183-1188 FEB 14 2008 2008

ITEM IDENTIFIER: doi:10.1038/sj.onc.1210722

ISSN: 0950-9232

DOCUMENT TYPE: Article; Editorial

RECORD TYPE: Abstract

LANGUAGE: English

ABSTRACT: I kappa B kinase (IKK) complex plays an important role in the regulation of signaling pathway that activates nuclear factor-kappa-B (NF-kappa B). Recently, we reported that cisplatin (CDDP) treatment causes a remarkable nuclear accumulation of $\text{IKK}\alpha$ in association with stabilization and activation of p73 . However, underlying mechanisms of CDDP-induced nuclear accumulation of $\text{IKK}\alpha$ are elusive. Here, we found that ataxia-telangiectasia mutated (ATM) is one of upstream mediators of $\text{IKK}\alpha$ during CDDP-induced apoptosis. In response to CDDP, ATM was phosphorylated at Ser-1981, which was accompanied with nuclear accumulation of $\text{IKK}\alpha$ in HepG2 cells, whereas CDDP treatment had undetectable effects on $\text{IKK}\alpha$ in ATM-deficient cells. Indirect immuno fluorescence experiments demonstrated that phosphorylated form of ATM colocalizes with nuclear $\text{IKK}\alpha$ in response to CDDP. In vitro kinase assay indicated that ATM phosphorylates $\text{IKK}\alpha$ at Ser-473. Moreover, $\text{IKK}\alpha$ -deficient MEFs displayed CDDP-resistant phenotype as compared with wild-type MEFs. Taken together, our present results suggest that ATM-mediated phosphorylation of nuclear $\text{IKK}\alpha$, which stabilizes p73 , is one of the main apoptotic pathways in response to CDDP.

1/7/2

DIALOG(R)File 5:Biosis Previews(R)

(c) 2009 The Thomson Corporation. All rts. reserv.

0019802300 BIOSIS NO.: 200700462041

Stabilization of p73^{Mdm2} by nuclear I kappa B kinase- α mediates cisplatin-induced apoptosis

AUTHOR: Furuya Kazushige; Ozaki Toshinori; Hanamoto Takayuki; Hosoda Mitsuchika; Hayashi Syunji; Barker Philip A; Takano Kunio; Matsumoto Masahiko; Nakagawara Akira (Reprint)

AUTHOR ADDRESS: Canc Ctr Res Inst, Div Biochem, Chuo Ku, 666-2 Nitona, Chiba 2608717, Japan**Japan

AUTHOR E-MAIL ADDRESS: akiranak@chiba-cc.jp

JOURNAL: Journal of Biological Chemistry 282 (25): p18365-18378 JUN 22 2007 2007

ITEM IDENTIFIER: doi:10.1074/jbc.M610522200

ISSN: 0021-9258

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

LANGUAGE: English

ABSTRACT: In response to DNA damage, p53 and its homolog p73^{Mdm2} have a function antagonistic to NF- κ B in deciding cell fate. Here, we show for the first time that p73^{Mdm2} , but not p53, is stabilized by physical interaction with nuclear I kappa B kinase ($\text{IKK}\alpha$)- α to enhance cisplatin (CDDP)-induced apoptosis. CDDP caused a significant increase in the amounts of nuclear $\text{IKK}\alpha$ - α and p73^{Mdm2} α in human osteosarcoma-derived U2OS cells. Ectopic expression of $\text{IKK}\alpha$ - α prolonged the half-life of p73^{Mdm2} by inhibiting its ubiquitination and thereby enhancing its transactivation and pro-apoptotic activities. Consistent with these results, small interfering RNA-mediated knockdown of endogenous $\text{IKK}\alpha$ - α inhibited the CDDP-mediated accumulation of p73^{Mdm2} α . The kinase-deficient mutant form of $\text{IKK}\alpha$ - α interacted with p73^{Mdm2} α , but failed to stabilize it. Furthermore, CDDP-mediated accumulation of endogenous p73^{Mdm2} α was not detected in mouse embryonic fibroblasts (MEFs) prepared from $\text{IKK}\alpha$ - α - α -deficient mice, and CDDP sensitivity was significantly decreased in $\text{IKK}\alpha$ - α -deficient MEFs compared with wild-type MEFs. Thus, our results strongly suggest that the nuclear $\text{IKK}\alpha$ - α -mediated accumulation of p73^{Mdm2} α is one of the novel molecular mechanisms to induce apoptotic cell death in response to CDDP, which may be particularly important in killing tumor cells with p53 mutation.

? e au=nakagawara akira

Ref	Items	Index-term
E1	1	AU=NAKAGAWAJI K
E2	109	AU=NAKAGAWARA A
E3	174	*AU=NAKAGAWARA AKIRA
E4	1	AU=NAKAGAWARA AKITA
E5	2	AU=NAKAGAWARA CHIHAYA
E6	3	AU=NAKAGAWARA EIKI
E7	37	AU=NAKAGAWARA G
E8	1	AU=NAKAGAWARA GIZIO
E9	27	AU=NAKAGAWARA GIZO
E10	8	AU=NAKAGAWARA GIZOU
E11	4	AU=NAKAGAWARA H
E12	4	AU=NAKAGAWARA HIROSHI

Enter P or PAGE for more

```
? s e3
      S2      174  AU='NAKAGAWARA AKIRA'
? s s2 and p73
      174  S2
      1565 P73
      S3      32  S2 AND P73
? s s2 and IKK?
      174  S2
      2942 IKK?
      S4      1  S2 AND IKK?
? t s4/7/1
```

4/7/1
 DIALOG(R)File 5:Biosis Previews(R)
 (c) 2009 The Thomson Corporation. All rts. reserv.

0019802300 BIOSIS NO.: 200700462041
 Stabilization of p73 by nuclear I kappa B kinase-alpha mediates
 cisplatin-induced apoptosis
 AUTHOR: Furuya Kazushige; Ozaki Toshinori; Hanamoto Takayuki; Hosoda
 Mitsuchika; Hayashi Syunji; Barker Philip A; Takano Kunio; Matsumoto
 Masahiko; %Nakagawara Akira% (Reprint)
 AUTHOR ADDRESS: Canc Ctr Res Inst, Div Biochem, Chuo Ku, 666-2 Nitona,
 Chiba 2608717, Japan**Japan
 AUTHOR E-MAIL ADDRESS: akiranak@chiba-cc.jp
 JOURNAL: Journal of Biological Chemistry 282 (25): p18365-18378 JUN 22
 2007 2007
 ITEM IDENTIFIER: doi:10.1074/jbc.M610522200
 ISSN: 0021-9258
 DOCUMENT TYPE: Article
 RECORD TYPE: Abstract
 LANGUAGE: English

ABSTRACT: In response to DNA damage, p53 and its homolog p73 have a
 function antagonistic to NF-kappa B in deciding cell fate. Here, we show
 for the first time that p73, but not p53, is stabilized by physical
 interaction with nuclear I kappa B kinase (IKK)-alpha to enhance
 cisplatin (CDDP)-induced apoptosis. CDDP caused a significant increase in
 the amounts of nuclear IKK-alpha and p73 alpha in human
 osteosarcoma-derived U2OS cells. Ectopic expression of IKK-alpha
 prolonged the half-life of p73 by inhibiting its ubiquitination and
 thereby enhancing its transactivation and pro-apoptotic activities.
 Consistent with these results, small interfering RNA-mediated knockdown
 of endogenous IKK-alpha inhibited the CDDP-mediated accumulation of
 p73 alpha. The kinase-deficient mutant form of IKK-alpha interacted
 with p73 alpha, but failed to stabilize it. Furthermore, CDDP-mediated
 accumulation of endogenous p73 alpha was not detected in mouse embryonic
 fibroblasts (MEFs) prepared from IKK-alpha-deficient mice, and CDDP
 sensitivity was significantly decreased in IKK-alpha-deficient MEFs
 compared with wild-type MEFs. Thus, our results strongly suggest that the
 nuclear IKK-alpha-mediated accumulation of p73 alpha is one of the
 novel molecular mechanisms to induce apoptotic cell death in response to
 CDDP, which may be particularly important in killing tumor cells with p53
 mutation.

? ds

Set	Items	Description
-----	-------	-------------

```
S1          2   P73 AND IKK?
S2         174  AU='NAKAGAWARA AKIRA'
S3          32  S2 AND P73
S4          1   S2 AND IKK?
? log y
    09feb09 10:15:43 User217744 Session D1185.3
      $6.58    1.063 DialUnits File5
        $7.32  3 Type(s) in Format 7
          $7.32  3 Types
    $13.90 Estimated cost File5
    $0.80 TELNET
    $14.70 Estimated cost this search
    $15.23 Estimated total session cost    1.447 DialUnits
Logoff: level 05.24.00 D 10:15:43
```